

MINIMAL AREA ELLIPSES IN THE HYPERBOLIC PLANE

MATTHIAS J. WEBER AND HANS-PETER SCHRÖCKER

ABSTRACT. We present uniqueness results for enclosing ellipses of minimal area in the hyperbolic plane. Uniqueness can be guaranteed if the minimizers are sought among all ellipses with prescribed axes or center. In the general case, we present a sufficient and easily verifiable criterion on the enclosed set that ensures uniqueness.

1. INTRODUCTION AND STATEMENT OF THE MAIN RESULT

By a well-known theorem of convex geometry, a full-dimensional, compact subset F of the Euclidean plane can be enclosed by a unique ellipse C of minimal area. We share the general belief that this is an important but easy result. Therefore, it is not surprising that recently much more general uniqueness results were obtained [8, 16, 17]. These articles also contain more complete references to the relevant literature.

The situation in the elliptic plane is different: As of today, uniqueness can only be guaranteed for “sufficiently small and round sets F ”. The precise statement can be found in [18, Theorem 8]. Its proof requires some non-trivial calculations. It is still an open question whether any compact subset of the elliptic plane possesses a unique enclosing conic or not.

In this article we consider uniqueness of the minimal area ellipse in the hyperbolic plane. The algebraic equivalence of elliptic and hyperbolic geometries suggests to imitate the proof of [18, Theorem 8]. Indeed, this is possible to a large extent, but not completely. The outcome of this research is similar to the elliptic case. Uniqueness can be guaranteed if some conditions on the axis lengths of enclosing ellipses of minimal area are met. If this is not possible, we can make neither a positive nor a negative uniqueness statement. Our main result is

Theorem 1. *Consider a compact and full-dimensional subset F of the hyperbolic plane. The enclosing ellipse of minimal area to F is unique if the following conditions are met:*

- *There exist positive numbers ϱ, R such that the semi-axis lengths of the (a priori not necessarily unique) minimal ellipses are in the closed interval $[\varrho, R]$.*
- *The values $\nu_1 = \coth^2 R$ and $\nu_2 = \coth^2 \varrho$ satisfy the inequality*

$$(1) \quad H(\nu_1, \nu_2) := -13\nu_1^2 + 5\nu_1\nu_2 - 3\nu_1 + 7\nu_2 + 4 \leq 0.$$

The minimal enclosing ellipse $C_{\min}$ to the convex hull F of a finite point set is depicted in Figure 2. The drawing refers to the Cayley-Klein model of the hyperbolic plane which will be introduced in Section 2.

Figure 1 depicts the graph of the function $H(\nu_1, \nu_2) = 0$. The shaded area contains admissible values ν_1, ν_2 . The meaning of the remaining elements will be explained later in the text.

Date: July 7, 2011.

2010 Mathematics Subject Classification. 52A40; 52A55, 51M10.

Key words and phrases. Hyperbolic geometry, enclosing ellipse, minimal area, uniqueness.




FIGURE 1. The curves $H(\nu_1, \nu_2) = 0$ and $h_2(\nu_1, \nu_2) = 0$.

Condition (1) is certainly fulfilled for $\nu_1 = \nu_2$ and $\nu_1 \rightarrow \infty$. Thus, Theorem 1 informally states that minimal enclosing ellipses are unique if they are sufficiently small and round. Of course, the Theorem should be accompanied by an easily verifiable criterion that ensure suitably shaped minimal ellipses:

Proposition 2. *Consider a compact and full-dimensional subset F of the hyperbolic plane and denote its (hyperbolic) convex hull by $\bar{F}$. Assume $\bar{F}$ admits an inscribed circle of radius ϱ and a circumscribed ellipse of area S . Denote by R the major semi-axis length of an ellipse of area S and minor semi-axis length r . Then the minimal area ellipse of F has semi-axis length in the interval $[\varrho, R]$.*

We omit the obvious proof of this proposition. Together with Theorem 1, it leads to the following sufficient test for the uniqueness of the minimal enclosing ellipsoid to a given set F :

- (1) Find a (large) inscribed circle to $\bar{F}$ and denote its radius by ϱ .
- (2) Find a (small) circumscribed ellipse to $\bar{F}$ and denote its area by S .
- (3) Compute the unique value R such that an ellipse with semi-axis lengths ϱ and R has area S . By construction, $[\varrho, R]$ is not empty.
- (4) The minimal area ellipse is unique, if ϱ and R satisfy the inequality (1).

We will show later that if a point (ν_1, ν_2) satisfies (1) then the same is true for every admissible point (ν'_1, ν'_2) with $\nu'_1 \geq \nu_1$ and $\nu'_2 \leq \nu_2$. This means, that chances for an affirmative uniqueness statement increase with a large value of the radius ϱ and a small value of the area S , that is, with the quality of the input obtained from the first and the second step.

The basic ideas and the initial calculations in our proof of Theorem 1 are more or less identical to the proof of [18, Theorem 8]. The minor differences pertain to occasional changes in sign and the use of the hyperbolic functions $\cosh$, $\sinh$, etc. instead of their spherical counterparts $\cos$, $\sin$, etc. The major differences are in the final estimates. Given the similarities between the elliptic and hyperbolic case, we consider a rather terse presentation appropriate. Yet, we will try to work out the crucial junctions points and the major differences.

In Section 2 we settle our notation and introduce the hyperboloid model of the hyperbolic plane, where our calculations take place. In Section 3 we provide a formula for the area of ellipses in the hyperbolic plane, which is probably hard to find elsewhere. In Section 4, we prove a fundamental convexity result for the area function. By standard arguments, it yields uniqueness of the minimal ellipse among all ellipses with prescribed axes or center. The proof of Theorem 1 is given

in Section 5. Its main ingredient is Lemma 7, the Half-Turn Lemma. The merely technical parts of its proof are moved to the appendix.

2. THE HYPERBOLOID MODEL OF HYPERBOLIC GEOMETRY

In [18] we used the spherical model of the elliptic plane for investigating uniqueness of minimal area conics. It is obtained from the geometry of the unit sphere $\mathbb{S}^2$ of Euclidean three-space by identifying antipodal points. By analogy, our calculations in this article refer to the spherical model (or “hyperboloid model”) of the hyperbolic plane which is obtained in similar fashion from the geometry of the sphere of squared radius -1 in Minkowski three space $\mathbb{R}_1^3$. An elementary introduction to this model is given in [14].

Minkowski three-space $\mathbb{R}_1^3$ is the metric space over $\mathbb{R}^3$ where the metric is induced by the indefinite inner product

$$\langle x, y \rangle = -x_0y_0 + x_1y_1 + x_2y_2.$$

The locus of the spherical model of the hyperbolic plane is the sphere $\mathbb{S}_1^2$, defined as

$$\mathbb{S}_1^2 = \{x \in \mathbb{R}_1^3 : \|x\|^2 = -x_0^2 + x_1^2 + x_2^2 = -1\}.$$

In a Euclidean interpretation, it is a hyperboloid of two sheets. We use $\mathbb{S}_1^2$ as a model of the hyperbolic plane $\mathbb{H}^2$. The following concepts are taken from [14]:

- The points of $\mathbb{H}^2$ are the points of $\mathbb{S}_1^2$ with antipodal points x and $-x$ identified.
- The lines of $\mathbb{H}^2$ are the intersections of $\mathbb{S}_1^2$ with planes through the origin 0.
- The hyperbolic distance between two points $x, y \in \mathbb{S}_1^2$ is defined by

$$\text{dist}(x, y) = \text{arccosh}(-\langle x, y \rangle).$$

- The hyperbolic angle between two straight lines K and L is defined by

$$\angle(k, l) = \arccos \frac{\langle k, l \rangle}{\|k\| \cdot \|l\|}$$

where k and l are two arbitrary tangent vectors of K and L , respectively.

Note that this model of $\mathbb{H}^2$ is closely related to the well-known bundle model and also the Cayley-Klein model of the hyperbolic plane. The bundle model is obtained by connecting points and lines from the spherical model with the origin 0 of $\mathbb{R}_1^3$; the Cayley-Klein model is obtained by intersecting the bundle model with the plane $x_0 = 1$. Its points are the inner points of the circle

$$K: x_0 = 1, x_1^2 + x_2^2 = 1.$$

We will occasionally use the Cayley-Klein model for the purpose of visualization but it is also convenient for defining center and axes of an ellipse C in the hyperbolic plane.

The conics in the spherical model of $\mathbb{H}^2$ are the intersections of $\mathbb{S}_1^2$ with quadratic cones centered at 0. In the Cayley-Klein model, hyperbolic ellipses are conics that lie in the interior of K . The ellipse center is the unique vertex c of the common polar triangle P of C and K . It is indeed a center in elementary sense, as it halves the (hyperbolic) distance between the ellipse points on any line incident with c . The axes of C are the two sides of P through c . Degenerate pole triangles characterize the circles among the ellipses. Their center is still well-defined but the axes are undetermined so that any line through c can be addressed as axis. Figure 2 displays a hyperbolic ellipse, its center and axes in the Cayley-Klein model.




FIGURE 2. Center c and axes of a hyperbolic ellipse C ; minimal ellipse $C_{\min}$ to the convex hull F of a finite points set

3. THE AREA OF ELLIPSES

The hyperbolic plane $\mathbb{H}^2$ can be parametrized as

$$(2) \quad \mathbb{H}^2: Y(\theta, \varphi) = \begin{pmatrix} \cosh \theta \\ \sinh \theta \sin \varphi \\ \sinh \theta \cos \varphi \end{pmatrix}, \quad \theta \in [0, \infty), \varphi \in [-\pi, \pi).$$

A conic C , defined as the intersection of this point set with a quadratic cone whose vertex is in the origin, can be described as

$$C = \{x \in \mathbb{H}^2: x^T \cdot M \cdot x = 0\},$$

where $M \in \mathbb{R}^{3 \times 3}$ is an indefinite symmetric matrix of full rank. A vector x is called (Minkowski) eigenvector of M with (Minkowski) eigenvalue λ if

$$(3) \quad M \cdot x = \lambda I \cdot x. \quad \text{where } I = \text{diag}(-1, 1, 1).$$

By $e(M) = (\nu_0, \nu_1, \nu_2)$ we denote the vector of eigenvalues of M , arranged in ascending order. We will only consider the case where M describes an ellipse. In this case M can be normalized such that $e(M) = (1, \nu_1, \nu_2)$ and $1 < \nu_1 \leq \nu_2$.

A point x is contained in the ellipse C if it satisfies $x^T \cdot M \cdot x < 0$ and M is in normal form. After a suitable (Minkowski) rotation of $\mathbb{S}_1^2$ we may assume that the ellipse is described by the diagonal matrix

$$(4) \quad M = \text{diag}(-1, \nu_1, \nu_2).$$

Referring to the parametrization (2), points inside C belong to parameter values (θ, φ) related by

$$\theta < \theta^* = \text{arccosh} \sqrt{\frac{\nu_1 \sin^2 \varphi + \nu_2 \cos^2 \varphi}{\nu_1 \sin^2 \varphi + \nu_2 \cos^2 \varphi - 1}}.$$

By integrating the area element

$$\sqrt{\left\langle \frac{\partial H}{\partial \theta}, \frac{\partial H}{\partial \theta} \right\rangle \cdot \left\langle \frac{\partial H}{\partial \varphi}, \frac{\partial H}{\partial \varphi} \right\rangle - \left\langle \frac{\partial H}{\partial \theta}, \frac{\partial H}{\partial \varphi} \right\rangle^2} d\theta \wedge d\varphi = \sinh \theta d\theta \wedge d\varphi$$

of (2) (see for example Proposition 5.2 of [3]) we obtain the area of the conic C as

$$\begin{aligned}
 \text{area}(C) &= \text{area}(\nu_1, \nu_2) = \int_{-\pi}^{\pi} \int_0^{\theta^*} \sinh \theta \, d\theta \, d\varphi \\
 (5) \quad &= \int_{-\pi}^{\pi} (\cosh \theta^* - 1) \, d\varphi = \int_{-\pi}^{\pi} \sqrt{\frac{\nu_1 \sin^2 \varphi + \nu_2 \cos^2 \varphi}{\nu_1 \sin^2 \varphi + \nu_2 \cos^2 \varphi - 1}} \, d\varphi - 2\pi.
 \end{aligned}$$

This is valid as long as M is normalized such that $e(M) = (1, \nu_1, \nu_2)$. If M is not normalized and has ordered eigenvalues $e(M) = (\nu_0, \nu_1, \nu_2)$, the area formula becomes

$$(6) \quad \text{area}(\nu_0, \nu_1, \nu_2) = \int_{-\pi}^{\pi} \sqrt{\frac{\nu_1 \sin^2 \varphi + \nu_2 \cos^2 \varphi}{\nu_1 \sin^2 \varphi + \nu_2 \cos^2 \varphi - \nu_0}} \, d\varphi - 2\pi.$$

4. CONVEXITY OF THE AREA FUNCTION.

Convexity of the area function (5) is already the key property for uniqueness of the minimal area ellipse among concentric or co-axial ellipses. Recall that only values $\nu_1, \nu_2 > 1$ are admissible.

Lemma 3. *The area function (5) is strictly convex for $\nu_1, \nu_2 > 1$.*

Proof. We proof that the Hessian matrix of (5) is positive definite, that is, all its principal minors are positive. The upper left entry equals

$$(7) \quad \frac{\partial^2 \text{area}}{\partial \nu_1^2} = \frac{1}{4} \int_{-\pi}^{\pi} J \sin^4 \varphi \, d\varphi,$$

where

$$J = \frac{4\nu_1 \sin^2 \varphi + 4\nu_2 \cos^2 \varphi - 1}{(\nu_1 \sin^2 \varphi + \nu_2 \cos^2 \varphi)^{3/2} (\nu_1 \sin^2 \varphi + \nu_2 \cos^2 \varphi - 1)^{5/2}}.$$

Clearly, J is positive for admissible values of ν_1 and ν_2 . Therefore, (7) is positive as well. The determinant of the Hessian matrix is

$$\begin{aligned}
 (8) \quad &\frac{\partial^2 \text{area}}{\partial \nu_1^2} \frac{\partial^2 \text{area}}{\partial \nu_2^2} - \left(\frac{\partial \text{area}}{\partial \nu_1 \partial \nu_2} \right)^2 \\
 &= \frac{1}{16} \int_{-\pi}^{\pi} J \sin^4 \varphi \, d\varphi \cdot \int_{-\pi}^{\pi} J \cos^4 \varphi \, d\varphi - \frac{1}{16} \left(\int_{-\pi}^{\pi} J \sin^2 \varphi \cos^2 \varphi \, d\varphi \right)^2.
 \end{aligned}$$

Because $\sqrt{J} \sin^2 \varphi$ and $\sqrt{J} \cos^2 \varphi$ are not proportional we can apply the strict Schwarz inequality and find

$$\sqrt{\int_{-\pi}^{\pi} (\sqrt{J} \sin^2 \varphi)^2 \, d\varphi} \cdot \sqrt{\int_{-\pi}^{\pi} (\sqrt{J} \cos^2 \varphi)^2 \, d\varphi} > \int_{-\pi}^{\pi} J \sin^2 \varphi \cos^2 \varphi \, d\varphi.$$

Thus, (8) is positive and $\text{area}(\nu_1, \nu_2)$ is indeed a strictly convex function. $\square$

Now, two uniqueness results follow from standard arguments (see [16, 17] and in particular [18]).

Theorem 4. *Let F be a compact and full-dimensional subset of the hyperbolic plane. Among all ellipses with two given axes that contain F there exists exactly one with minimal area.*

Theorem 5. *Let F be a compact and full-dimensional subset of the hyperbolic plane. Among all ellipses with given center that contain F there exists exactly one with minimal area.*

We give a quick outline of the proofs of Theorem 4 and 5, mainly because this gives us the opportunity to introduce an important concept that will be required later.

Definition 6 (in-between ellipse). Let C_0 and C_1 be two ellipses

$$C_i = \{x \in \mathbb{H}^2 : x^T \cdot M_i \cdot x = 0\}, \quad i = 0, 1$$

where the matrices M_i are indefinite and have Minkowski eigenvalues $\nu_{i,0} = 1$ and $\nu_{i,1}, \nu_{i,2} > 1$. For $\lambda \in (0, 1)$, the *in-between ellipse* C_λ of C_0 and C_1 is defined as

$$C_\lambda = \{x \in \mathbb{H}^2 : x^T \cdot M_\lambda \cdot x\},$$

where

$$M_\lambda = (1 - \lambda)M_0 + \lambda M_1.$$

We also write $C_\lambda = (1 - \lambda)C_0 + \lambda C_1$.

It is obvious that C_λ contains the common interior of C_0 and C_1 and is an ellipse if this interior is not empty. Moreover, it follows from Lemma 3 and the strict version of Davis' convexity theorem [4, 11] that $\text{area}(C_\lambda)$ is a strictly convex function of λ . More detailed arguments can be found in [17, 18]. The important fact to remember is that two enclosing conics C_0 and C_1 of the same area give rise to an in-between conic C_λ of lesser area. Thus, the assumption of two minimal area conics leads to a contradiction. Note that convexity of $\text{area}(C_\lambda)$ in the general (non-concentric) case is not implied by Davis' convexity theorem.

5. UNIQUENESS IN THE GENERAL CASE.

Now we come to the proof of Theorem 1, the general uniqueness result. As usual, existence follows from compactness arguments. The basic ideas and initial steps in the proof of uniqueness are not different from the proof of Theorem 8 in [18]. We give an outline:

- Assume existence of two minimal enclosing ellipses C_0 and C_1 .
- Find the unique (hyperbolic) half-turn η (an idempotent hyperbolic rotation) such that $C_1^* = \eta(C_1)$ and C_1^* is concentric with C_0 .
- Define in-between ellipses $C_\lambda = (1 - \lambda)C_0 + \lambda C_1$ and $C_\lambda^* = (1 - \lambda)C_0 + \lambda C_1^*$ according to Definition 6.
- Show that there exists $\varepsilon > 0$ such that $\text{area}(C_\lambda) < \text{area}(C_\lambda^*)$ for $0 < \lambda < \varepsilon$. Because of $\text{area}(C_\lambda^*) \leq \text{area}(C_0) = \text{area}(C_1)$ (with equality iff $C_1^* = C_0$) this contradicts the assumed minimality of C_0 and C_1 .

Existence of ε in the last step of this program can be proved by showing the inequality

$$(9) \quad \left. \frac{\partial \text{area}(C_\lambda^*)}{\partial \lambda} \right|_{\lambda=0} < \left. \frac{\partial \text{area}(C_\lambda)}{\partial \lambda} \right|_{\lambda=0}.$$

The advantage of this approach is that both sides of (9) can be readily computed from the normalized equations that describe C_0 , C_1 , and C_1^* . In particular, the cubic problem of calculating the eigenvalues of the matrices describing C_λ or C_λ^* is avoided.

In order to follow the outline of the proof of Theorem 1 we have to compute the ellipses C_0 , C_1 and C_1^* in a sufficiently general way. By Theorem 5, the centers of C_0 and C_1 can be assumed to be different. Thus, there exists a unique mid-point r of their respective centers c_0 and c_1 . Define C_1^* as the ellipse obtained by applying the half-turn with center r to C_1 . The ellipses C_0 , C_1 , and C_1^* are described by matrices M_0 , M_1 , and M_1^* with respective eigenvalues

$$e(M_0) = (1, \nu_{0,1}, \nu_{0,2}), \quad e(M_1) = e(M_1^*) = (1, \nu_{1,1}, \nu_{1,2}).$$




FIGURE 3. The case of two circles

We would like to make some admissible assumptions on these eigenvalues. Because of $\text{area}(C_0) = \text{area}(C_1)$, we have

$$(10) \quad 1 < \nu_{0,1} \leq \nu_{1,1} \leq \nu_{1,2} \leq \nu_{0,2}.$$

If $\nu_{0,1} = \nu_{1,1}$ or $\nu_{1,2} = \nu_{0,2}$, (10) holds with equality throughout and both ellipses are actually congruent circles. In this case a simple construction produces a smaller enclosing circle (Figure 3): Denote the two intersection points of C_0 and C_1 by s_0 and s_1 . By elementary hyperbolic geometry, the circle S over the diameter s_0, s_1 is smaller than C_0 and C_1 and it contains the common interior of C_0 and C_1 .

Thus, the case of two congruent circles can be excluded and we may assume that the eigenvalues of M_0 and M_1 are ordered according to

$$(11) \quad 1 < \nu_{0,1} < \nu_{1,1} \leq \nu_{1,2} < \nu_{0,2}.$$

Now we compute the derivative of the area function (6) with respect to λ . For that purpose, we assume that C_0 is given by the normal form (4) and C_1 is obtained from an ellipse in this normal form by a hyperbolic rotation, that is,

$$(12) \quad M_0 = \text{diag}(-1, \nu_{0,1}, \nu_{0,2}), \quad M_1 = (Q^{-1})^T \cdot \text{diag}(-1, \nu_{1,1}, \nu_{1,2}) \cdot Q^{-1}$$

with the hyperbolic rotation matrix

$$(13) \quad Q = \begin{pmatrix} q_0^2 + q_1^2 + q_2^2 + q_3^2 & 2(q_0q_3 + q_1q_2) & 2(q_1q_3 - q_0q_2) \\ 2(q_0q_3 - q_1q_2) & q_0^2 - q_1^2 - q_2^2 + q_3^2 & 2(q_0q_1 - q_2q_3) \\ 2(-q_0q_2 - q_1q_3) & 2(-q_0q_1 - q_2q_3) & q_0^2 - q_1^2 + q_2^2 - q_3^2 \end{pmatrix},$$

$$q_0^2 + q_1^2 - q_2^2 - q_3^2 = 1.$$

Up to the irrelevant sign of q_1 and an index shift, this is precisely Equation (5) of [13]. The rotation angle -2ξ is given by $q_0 = \cos \xi$, the axis direction is $(-q_1, q_2, q_3)^T$. A hyperbolic half-turn is obtained by substituting $q_0 = 0$ into (13). In this case, $Q \cdot Q$ indeed equals the unit matrix $\text{diag}(1, 1, 1)$.

The matrix M_λ to C_λ is computed according to Definition 6. Its ordered eigenvalues (ν_0, ν_1, ν_2) are functions of λ . In the vicinity of $\lambda = 0$ we have $\nu_0(\lambda) > 0$ and $1 < \nu_1(\lambda) < \nu_2(\lambda)$. These eigenvalues are implicitly defined as roots of the characteristic polynomial $P(\lambda, \nu(\lambda)) = \det(M_\lambda - \nu I)$ of M_λ where I is the matrix defined in (3). For $\lambda = 0$ we know the values of these roots:

$$\nu_0(0) = 1, \quad \nu_1(0) = \nu_{0,1}, \quad \nu_2(0) = \nu_{0,2}.$$

By implicit derivation we have

$$\frac{d\nu_i}{d\lambda}(0) = -\frac{\frac{\partial P}{\partial \lambda}(0, \nu_i(0))}{\frac{\partial P}{\partial \nu}(0, \nu_i(0))}, \quad i = 0, 1, 2.$$

Furthermore, we can compute the partial derivatives

$$\frac{\partial \text{area}(\nu_0, \nu_1, \nu_2)}{\partial \nu_i}, \quad i = 0, 1, 2$$

of (6). Using the chain rule

$$\frac{\partial \text{area}(C_\lambda)}{\partial \lambda} \Big|_{\lambda=0} = \frac{\partial \text{area}}{\partial \nu_0} \frac{\partial \nu_0}{\partial \lambda} \Big|_{\lambda=0} + \frac{\partial \text{area}}{\partial \nu_1} \frac{\partial \nu_1}{\partial \lambda} \Big|_{\lambda=0} + \frac{\partial \text{area}}{\partial \nu_2} \frac{\partial \nu_2}{\partial \lambda} \Big|_{\lambda=0},$$

we find

$$(14) \quad \frac{\partial \text{area}(C_\lambda)}{\partial \lambda} \Big|_{\lambda=0} = -\frac{1}{2} \int_{-\pi}^{\pi} \frac{D}{N} d\varphi$$

where

$$\begin{aligned} D &= ((q_{1,2}^2 \nu_{1,2} + q_{1,3}^2 \nu_{1,2} - q_{1,1}^2) \nu_{0,1} + q_{2,2}^2 \nu_{1,1} + q_{2,3}^2 \nu_{1,2} - q_{2,1}^2) \sin^2 \varphi + \\ &\quad ((q_{1,2}^2 \nu_{1,1} + q_{1,3}^2 \nu_{1,2} - q_{1,1}^2) \nu_{0,2} + q_{3,2}^2 \nu_{1,1} + q_{3,3}^2 \nu_{1,2} - q_{3,1}^2) \cos^2 \varphi, \\ N &= (\nu_{0,1} \sin^2 \varphi + \nu_{0,2} \cos^2 \varphi - 1)^{3/2} (\nu_{0,1} \sin^2 \varphi + \nu_{0,2} \cos^2 \varphi)^{1/2} \end{aligned}$$

and $q_{i,j}$ are the entries of the matrix (13). It will be convenient to write (14) in terms of the first and second complete elliptic integrals

$$(15) \quad K(z) = \int_0^1 \frac{1}{\sqrt{1-t^2} \sqrt{1-z^2 t^2}} dt \quad \text{and} \quad E(z) = \int_0^1 \frac{\sqrt{1-z^2 t^2}}{\sqrt{1-t^2}} dt.$$

Since we will evaluate them only at

$$f = \sqrt{\frac{\nu_{0,2} - \nu_{0,1}}{(\nu_{0,2} - 1)\nu_{0,1}}},$$

we use the abbreviations $\bar{E} := E(f)$ and $\bar{K} := K(f)$. By (11), f is always real and between 0 and 1. Substituting

$$\varphi = \arcsin \sqrt{\frac{\nu_{0,2} - x}{\nu_{0,2} - \nu_{0,1}}},$$

and noting that $2(\nu_{0,2} - x)(x - \nu_{0,1}) = 2(\nu_{0,1} - \nu_{0,2}) \cos \varphi \sin \varphi$ we can express the derivative of the area function in terms of $\bar{E}$ and $\bar{K}$:

$$(16) \quad \frac{\partial \text{area}(C_\lambda)}{\partial \lambda} \Big|_{\lambda=0} = \frac{2}{\sqrt{(\nu_{0,2} - 1)\nu_{0,1}(\nu_{0,2} - \nu_{0,1})(\nu_{0,1} - 1)}} \\ (A(q_{1,2}^2 \nu_{1,1} + q_{1,3}^2 \nu_{1,2} - q_{1,1}^2) + B(q_{2,2}^2 \nu_{1,1} + q_{2,3}^2 \nu_{1,2} - q_{2,1}^2) + \Gamma(q_{3,2}^2 \nu_{1,1} + q_{3,3}^2 \nu_{1,2} - q_{3,1}^2))$$

where

$$(17) \quad \begin{aligned} A &= -\nu_{0,1}(\nu_{0,2} - \nu_{0,1})\bar{E}, \quad B = \nu_{0,2}(\nu_{0,1} - 1)\bar{K} - \nu_{0,1}(\nu_{0,2} - 1)\bar{E}, \\ \Gamma &= \nu_{0,1}(\nu_{0,1} - 1)(\bar{E} - \bar{K}). \end{aligned}$$

Having computed (16), the preparatory work for the final (big) step in the proof of Theorem 1 is completed. We formulate the last step as a lemma:

Lemma 7 (Hyperbolic Half-Turn Lemma). *Consider three ellipses C_0 , C_1 , C_1^* of equal area. Assume that*

- C_0 and C_1^* are concentric,
- C_1 is obtained from C_1^* by a half-turn,

- the eigenvalues $\nu_{i,1}$, $\nu_{i,2}$ of the normalized matrix M_i to C_i ($i = 0, 1$) satisfy (11), and
- $H(\nu_{i,1}, \nu_{i,2}) \leq 0$ where H is defined in Equation (1).

Then the area of $C_\lambda = (1 - \lambda)C_0 + \lambda C_1$ is smaller than the area of $C_\lambda^* = (1 - \lambda)C_0 + \lambda C_1^*$, at least in the vicinity of $\lambda = 0$.

In order to proof Lemma 7, we compare the derivatives of the areas of C_λ and C_λ^* with respect to λ at $\lambda = 0$. The ellipse C_1^* can be obtained from an ellipse in normal form (4) by a rotation about $(1, 0, 0)^T$ through ζ . We can compute the matrix M_1 as in (12) by substituting

$$q_0 = \cos \frac{\zeta}{2}, \quad q_1 = -\sin \frac{\zeta}{2}, \quad q_2 = q_3 = 0$$

into the matrix (13). Plugging this into Equation (16) yields

$$\frac{1}{2} \frac{\partial \text{area}(C_\lambda^*)}{\partial \lambda} \Big|_{\lambda=0} = \frac{D_1^*}{N_1^*}$$

where

$$D_1^* = -A + (B \cos^2 \zeta + \Gamma \sin^2 \zeta) \nu_{1,1} + (B \sin^2 \zeta + \Gamma \cos^2 \zeta) \nu_{1,2},$$

$$N_1^* = \sqrt{(\nu_{0,2} - 1) \nu_{0,1} (\nu_{0,2} - \nu_{0,1}) (\nu_{0,1} - 1)},$$

and A, B, Γ are as in (17).

The ellipse C_1 is obtained by a half-turn from C_1^* about the rotation axis defined by the unit vector $r = (r_1, r_2, r_3)^T$. The matrix Q in (13) is the product of the rotation matrix about $(1, 0, 0)^T$ through ζ and a half-turn rotation matrix about the unit vector r . The later is obtained by substituting

$$q_0 = 0, \quad q_1 = -r_1, \quad q_2 = r_2, \quad q_3 = r_3$$

into Equation (13). Plugging the entries of the product matrix into (16) yields

$$\frac{1}{2} \frac{\partial \text{area}(C_1)}{\partial \lambda} \Big|_{\lambda=0} \equiv \frac{D_1}{N_1} \pmod{(r_1^2 - r_2^2 - r_3^2 - 1)}$$

where $N_1 = N_1^*$ and

$$D_1 = 4r_2r_3((2A + B + \Gamma)r_1^2 + (B - \Gamma)r_2^2 - (B - \Gamma)r_3^2)(\nu_{1,1} - \nu_{1,2}) \sin \zeta \cos \zeta$$

$$+ (4(A + B)r_1^2r_2^2 - 4(A + \Gamma)r_1^2r_3^2 - 8(B - \Gamma)r_2^2r_3^2 + B - \Gamma)(\nu_{1,1} - \nu_{1,2}) \cos^2 \zeta$$

$$+ 4(A + B)r_1^2r_2^2(\nu_{1,2} - 1) + 4(A + \Gamma)r_1^2r_3^2(\nu_{1,1} - 1)$$

$$+ 4(B - \Gamma)r_2^2r_3^2(\nu_{1,1} - \nu_{1,2}) + \Gamma\nu_{1,1} + B\nu_{1,2} - A.$$

Now we are going to prove the inequality $D_1 - D_1^* < 0$ for $\zeta \in [0, \frac{\pi}{2}]$. We substitute $\zeta = 2 \arctan t$ into its left-hand side and obtain a rational expression in t . Clearing the positive denominator $(1 + t^2)^2$, we are left with a polynomial $P(t)$ of degree four whose negativity on $[0, 1]$ has to be shown. To do this, we write $P(t)$ with respect to the Bernstein basis as

$$(18) \quad P(t) = \sum_{i=0}^4 p_i B_i^4(t) \quad \text{where} \quad B_i^4(t) = \binom{4}{i} (1-t)^{4-i} t^i$$

and show non-positivity of the coefficients p_1, p_2, p_3 and negativity of the coefficients p_0 and p_4 . After a straightforward basis transformation and reducing modulo $r_1^2 - r_2^2 - r_3^2 - 1$ we find

$$p_0 = 4((\nu_{1,1} - 1)(A + B)r_1^2r_2^2 + (\nu_{1,2} - 1)(A + \Gamma)r_1^2r_3^2 + (\nu_{1,2} - \nu_{1,1})(B - \Gamma)r_2^2r_3^2),$$

$$p_1 = 4((\nu_{1,1} - 1)(A + B)r_1^2r_2^2 + (\nu_{1,2} - 1)(A + \Gamma)r_1^2r_3^2 + (\nu_{1,2} - \nu_{1,1})(B - \Gamma)r_2^2r_3^2)$$

$$+ 2r_2r_3(\nu_{1,2} - \nu_{1,1})(-(2A + B + \Gamma)r_1^2 - (B - \Gamma)r_2^2 + (B - \Gamma)r_3^2),$$

$$3p_2 = 8(\nu_{1,1} + \nu_{1,2} - 2)((A + B)r_1^2 r_2^2 + (A + \Gamma)r_1^2 r_3^2) \\ + 12r_2 r_3(\nu_{1,2} - \nu_{1,1})(-(2A + B + \Gamma)r_1^2 - (B - \Gamma)r_2^2 + (B - \Gamma)r_3^2),$$

$$p_3 = 8((\nu_{1,2} - 1)(A + B)r_1^2 r_2^2 + (\nu_{1,1} - 1)(A + \Gamma)r_1^2 r_3^2 - (\nu_{1,2} - \nu_{1,1})(B - \Gamma)r_2^2 r_3^2) \\ + 4r_2 r_3(\nu_{1,2} - \nu_{1,1})(-(2A + B + \Gamma)r_1^2 - (B - \Gamma)r_2^2 + (B - \Gamma)r_3^2),$$

$$p_4 = 16((\nu_{1,2} - 1)(A + B)r_1^2 r_2^2 + (\nu_{1,1} - 1)(A + \Gamma)r_1^2 r_3^2 - (\nu_{1,2} - \nu_{1,1})(B - \Gamma)r_2^2 r_3^2).$$

Recall now Equation (11) ($1 < \nu_{0,1} < \nu_{1,1} \leq \nu_{1,2} < \nu_{0,2}$) and $r_1^2 - r_2^2 - r_3^2 = 1$ and observe that

- $A < B < \Gamma$; this is proved in Lemma 9 and Lemma 10 in the appendix.
- $\Gamma < 0$; this follows from $\bar{E} < \bar{K}$ and $\nu_{0,1} > 1$.

Under these conditions, the negativity of p_0 is clear except when $r_2 = r_3 = 0$. But this is the concentric case $C_1 = C_1^*$ and need not be considered. The non-positivity of the coefficients p_1, p_2, p_3 , and the negativity of p_4 is shown in Lemmas 12 and 13 below. This concludes the proof of the Half-Turn Lemma and, thus, also the proof of Theorem 1.

Example 8. We use the prerequisites of Theorem 1 on the radii r and R in Lemma 12. But one might wonder whether the Half-Turn Lemma remains true without these assumptions. The answer to this question is negative. We can provide an example, where the polynomial $P(t)$ attains positive values on $(0, 1)$.

Substituting $r_1^2 = r_2^2 + r_3^2 + 1$, the coefficient p_1 can be written as

$$p_1 = \nu_{1,2}(4r_2^2 r_3(A + B)(r_3 - r_2) + 4r_3^2(A + \Gamma)(r_3^2 - r_2 r_3 + 1) - 2(2A + B + \Gamma)r_2 r_3) \\ + \nu_{1,1}(4(A + B)r_2^2(r_2^2 + r_2 r_3 + 1) + 4(A + \Gamma)r_2 r_3^2(r_2 + r_3) + 2(2A + B + \Gamma)r_2 r_3) \\ - 4(1 + r_2^2 + r_3^2)(r_2^2(A + B) + r_3^2(A + \Gamma)).$$

Assuming $r_2, r_3 > 0$ we see that

- the coefficient of $\nu_{1,1}$ is always negative and
- it is possible to choose $\nu_{0,1}, \nu_{0,2}, r_2$, and r_3 so that the coefficient of $\nu_{1,2}$ is positive.

Consequently p_1 can be made positive for large $\nu_{1,2}$. The choice

$$\nu_{0,1} = \nu_{1,1} = 1.1, \quad \nu_{0,2} = \nu_{1,2} = 90, \quad r_2 = 0.9 \cos(0.3), \quad r_3 = 0.9 \sin(0.3),$$

accomplishes this and even makes $P(t)$ attain positive values for $t \in (0, 1)$ (the zeros are $t \approx 0.1272$ and $t \approx 0.1389$). Note that this does not imply

$$\left. \frac{\partial \text{area}(C_\lambda)}{\partial \lambda} \right|_{\lambda=0} > 0$$

and, thus, constitutes no counter-example to the statement that the area of C_λ is smaller than the area of C_0 and C_1 . We are not aware of such a counter-example.

6. CONCLUSION AND FUTURE RESEARCH

We proved uniqueness results for minimal enclosing ellipses in the hyperbolic plane. The general result (Theorem 1) involves rather cumbersome but straightforward calculations. The differences to the elliptic case are mainly in the final estimates for the coefficients of the polynomial $P(t)$ in (18) and can be found in the appendix.

It is apparent that Theorem 1 leaves room for improvements. Pushing back the frontier dictated by the inequality (1) in Theorem 1 would be nice. Substantial steps towards answering the question whether the minimal area ellipse to all compact and full-dimensional sets F in the hyperbolic plane is unique or not would be great.

Note that there is a subtle difference to the situation in the elliptic plane. In [18], we presented an example from which we inferred that uniqueness in the elliptic

plane cannot be proved by means of our construction of in-between conics. In the hyperbolic plane, we are not aware of such a configuration. Example 8 only shows that the estimate of the derivative of the area function is insufficient. Thus, there is a certain hope that a general uniqueness result can be proved by means of our construction.

Since uniqueness or non-uniqueness of minimal enclosing ellipses in the elliptic and hyperbolic plane remains a difficult topic, one might try to aim at a weaker result and consider only “typical” (in the sense of Baire categories, see [6, 7]) convex sets $\bar{F}$.

We would also like to mention that [18] and this article are the only results on extremal quadrics in non-Euclidean geometries that we are aware of. We can conceive numerous possibilities for generalizations. They pertain to the dimension of the surrounding space, the type of the quadric and enclosed set (for example minimal enclosing hyperbolas to line sets as in [15]) the measure for the quadric’s size (volume, surface area etc.), and the replacement of “minimal enclosing” by “maximal inscribed” quadrics.

The attentive reader will have noticed that our method of proving Theorem 1 can be adapted to these generalizations. Having defined an “in-between” quadric Q_λ by means of a suitable matrix convex combination, it might be infeasible to compute the size Q_λ in a form that allows further processing. But, provided the size function’s derivatives with respect to the matrix eigenvalues can be computed, it is, at least in principle, possible to obtain an explicit formula for the derivative of the size of Q_λ for $\lambda = 0$. Its negativity has to be shown so that the uniqueness problem is made accessible to numerous tools and techniques related to inequalities.

In the Euclidean setting, the mere uniqueness result is less important than John’s characterization of it via his famous decomposition of the identity. The original reference is the old paper [10]. But, following [1], many contemporary authors considered this topic [2, 5, 8, 9, 12]. Elliptic and hyperbolic versions of John’s characterization seem to be a worthy topic of future research.

APPENDIX. PROOFS OF AUXILIARY RESULTS

Lemma 9. *For A and B as in (17) we have $A < B$.*

Proof. We show that $A - B < 0$. By (17) we have

$$A - B = (\nu_{0,1} - 1)(\nu_{0,1}\bar{E} - \nu_{0,2}\bar{K}).$$

This is negative because of $\bar{E} < \bar{K}$ and $1 < \nu_{0,1} < \nu_{0,2}$. □

Lemma 10. *For B and Γ as in (17) we have $B < \Gamma$.*

Proof. We let $\Delta = B - \Gamma$ and view Δ as a function of $\nu_{0,1}$ and $\nu_{0,2}$. Its negativity for $1 < \nu_{0,1} < \nu_{0,2}$ follows from three facts:

- $\Delta = 0$ for $\nu_{0,1} = \nu_{0,2}$ (this is obvious because in this case we have $\bar{E} = \bar{K}$),
- $\frac{\partial \Delta}{\partial \nu_{0,2}} = 0$ for $\nu_{0,1} = \nu_{0,2}$, and
- Δ is concave in $\nu_{0,2}$ for $1 < \nu_{0,1} < \nu_{0,2}$.

We compute the first partial derivative of Δ with respect to $\nu_{0,2}$:

$$\frac{\partial \Delta}{\partial \nu_{0,2}} = \frac{(2\nu_{0,2}(1 - \nu_{0,2}) + \nu_{0,1} - 1)\nu_{0,1}\bar{E}}{2\nu_{0,2}(\nu_{0,2} - 1)} + \frac{(2\nu_{0,2} - 1)(\nu_{0,1} - 1)\bar{K}}{2(\nu_{0,2} - 1)}.$$

It vanishes for $\nu_{0,1} = \nu_{0,2}$. The second partial derivative of Δ with respect to $\nu_{0,2}$ equals

$$\frac{\partial^2 \Delta}{\partial \nu_{0,2}^2} = \frac{(\nu_{0,1} - 1)}{4\nu_{0,2}^2(\nu_{0,2} - 1)^2} J_1 \quad \text{where} \quad J_1 = \nu_{0,2}(\nu_{0,1} - 1)\bar{K} - \nu_{0,1}(5\nu_{0,2} - 2)\bar{E}.$$

We have to show that it is negative. The factor before J_1 is positive. To see the negativity of J_1 itself we write it in the integral form (see (15))

$$J_1 = \int_0^1 \frac{J_2}{\sqrt{1-t^2}\sqrt{1-f^2t^2}} dt,$$

where

$$J_2 = \nu_{0,2}(\nu_{0,1} - 1) - \nu_{0,1}(5\nu_{0,2} - 2) \left(1 - t^2 \frac{\nu_{0,2} - \nu_{0,1}}{\nu_{0,1}(\nu_{0,2} - 1)}\right).$$

The term J_2 is linear in t^2 . For $t = 0$ it equals $2\nu_{0,1}(1 - 2\nu_{0,2}) - \nu_{0,2} < 0$ and for $t = 1$ it equals $-\nu_{0,2}(\nu_{0,1} - 1)(4\nu_{0,2} - 1)/(\nu_{0,2} - 1) < 0$. Thus, $J_2 < 0$ for $t \in [0, 1]$. This implies $J_1 < 0$ and we see that Δ is indeed concave for $1 < \nu_{0,1} < \nu_{0,2}$. $\square$

We will deduce non-positivity of the Bernstein coefficients $p_1, \dots, p_3$ from the inequality (1) and the additional inequalities

$$(19) \quad h_1(\nu_1, \nu_2) := \nu_2 - 5\nu_1 + 4 \leq 0,$$

$$(20) \quad h_2(\nu_1, \nu_2) := -5\nu_1^2 + \nu_1\nu_2 + \nu_1 + \nu_2 + 2 \leq 0,$$

$$(21) \quad h_3(\nu_1, \nu_2) := \nu_2^2 - 5\nu_1\nu_2 - 2\nu_1 + 4\nu_2 + 2 \leq 0,$$

$$(22) \quad h_4(\nu_1, \nu_2) := 5\nu_2^2 - 13\nu_1\nu_2 - 2\nu_1 + 6\nu_2 + 4 \leq 0,$$

$$(23) \quad h_5(\nu_1, \nu_2) := -5\nu_1^2 + \nu_1\nu_2 - \nu_1 + 3\nu_2 + 2 \leq 0,$$

$$(24) \quad h_6(\nu_1, \nu_2) := \nu_2^2 - 5\nu_1\nu_2 + 2\nu_2 + 2 \leq 0,$$

which are all simple consequences of (1). We state this in Lemma 11, below.

The assumptions of Theorem 1 guarantee that these inequalities are fulfilled for $\nu_1 = \nu_{0,1}$, $\nu_2 = \nu_{0,2}$. Thus, we only have to show that the inequalities are satisfied on the set

$$U := \{(\nu_1, \nu_2) \mid 1 < \nu_1 < \nu_2\}.$$

Lemma 11. (a) *If a point $(\nu_1, \nu_2) \in U$ satisfies the inequality (1), it also satisfies the inequalities (19)–(24).*

(b) *If a point (ν_1^*, ν_2^*) satisfies the inequalities (1), and (19)–(24), the same is true for all points $(\nu_1, \nu_2) \in U$ with $\nu_1 \geq \nu_1^*$ and $\nu_2 \leq \nu_2^*$.*

Proof. It is an elementary exercise to verify that the curves defined by the implicit equations H and h_j over the closure of U are the graphs of strictly monotone increasing functions of $F(\nu_1)$ and $f_j(\nu_1)$

$$F, f_j : [1, \infty) \rightarrow [1, \infty),$$

for $j \in \{1, \dots, 6\}$. This implies assertion (b).

The functions $f_j - F$ are strictly monotone increasing as well. This and the observation $F(1) = f_j(1) = 1$ implies assertion (a). $\square$

Figure 1 displays the hyperbolas $H(\nu_1, \nu_2) = 0$ and, as an example, $h_2(\nu_1, \nu_2) = 0$ together with the line $\nu_1 = \nu_2$. The remaining curves are depicted in light-gray. The region U is dotted.

Lemma 12. *The coefficients p_1 , p_2 , and p_3 are not positive.*

Proof. We substitute $r_1^2 = 1 + r_2^2 + r_3^2$ into p_1 and observe that $p_1 = 0$ and $\frac{\partial p_1}{\partial r_2} = 0$ if $r_2 = r_3 = 0$. The lemma's claim holds true if we can show that the r_2 -parameter lines of p_1 , viewed as a function of r_2 and r_3 , are strictly concave, that is,

$$(25) \quad \frac{\partial^2 p_1}{\partial r_2^2} = 8(A + B)(\nu_{1,1} - 1)(6r_2^2 + 1) + 8((A + B)\nu_{1,2} + (A + \Gamma)\nu_{1,1} - (2A + B + \Gamma))r_3^2 - 24(\nu_{1,2} - \nu_{1,1})(A + B)r_2r_3 < 0.$$

The coefficient of $r_2 r_3$ is positive, the remaining terms are negative. By the inequality of arithmetic and geometric means we have $r_2 r_3 \leq (r_2^2 + r_3^2)/2$. We insert this into (25) to obtain

$$(26) \quad \frac{\partial^2 p_1}{\partial r_2^2} \leq 4(A+B)(3(-\nu_{1,2} + 5\nu_{1,1} - 4)r_2^2 + 2(\nu_{1,1} - 1)) \\ + 4(-(A+B)\nu_{1,2} + (5A+3B+2\Gamma)\nu_{1,1} - 2(2A+B+\Gamma))r_3^2 < 0.$$

The first term is negative if $\nu_{1,2} - 5\nu_{1,1} + 4 \leq 0$. This is implied by $\nu_{0,2} - 5\nu_{0,1} + 4 \leq 0$ and thus follows from (19). In the second term the coefficient of r_3^2 needs closer investigation. We want to show its negativity. By (11) we have

$$(27) \quad (5A+3B+2\Gamma)\nu_{1,1} - (A+B)\nu_{1,2} - 2(2A+B+\Gamma) \leq \\ (5A+3B+2\Gamma)\nu_{0,1} - (A+B)\nu_{0,2} - 2(2A+B+\Gamma).$$

Using (15) and (17), we write the term on the right in its integral form:

$$(28) \quad (5A+3B+2\Gamma)\nu_{0,1} - (A+B)\nu_{0,2} - 2(2A+B+\Gamma) \\ = (\nu_{0,2} - \nu_{0,1}) \int_0^1 \frac{J_3}{\sqrt{1-t^2}\sqrt{1-f^2t^2}}$$

where

$$J_3 = -\frac{\nu_{0,2} - \nu_{0,1}}{\nu_{0,2} - 1}(2\nu_{0,2} - 7\nu_{0,1} + 5)t^2 + \nu_{0,2}(\nu_{0,1} + 1) + \nu_{0,1}(1 - 5\nu_{0,1}) + 2.$$

We see that J_3 is linear in t^2 . For $t = 0$ and $t = 1$ it attains the respective values

$$(29) \quad J_3|_{t=0} = -5\nu_{0,1}^2 + \nu_{0,1}\nu_{0,2} + \nu_{0,1} + \nu_{0,2} + 2,$$

$$(30) \quad J_3|_{t=1} = \frac{\nu_{0,1} - 1}{\nu_{0,2} - 1}(\nu_{0,2}^2 - 5\nu_{0,1}\nu_{0,2} - 2\nu_{0,1} + 4\nu_{0,2} + 2).$$

The right-hand side of (29) is not positive by (20). The right-hand side of (30) is not positive by (21). We conclude that the integrand J_3 is not positive for $t \in [0, 1]$ and the same is true for $\frac{\partial^2 p_1}{\partial r_2^2}$. Hence, the coefficient p_1 as a function of r_2 is concave with the maximum, $p_1 = 0$ attained at $r_2 = r_3 = 0$. Thus, p_1 is not positive.

The proofs of non-positivity of p_2 and p_3 run along exactly the same lines. We only provide the relevant formulas and reduce the explanatory text between them to a minimum. Equations (25) and (26) become

$$3 \frac{\partial^2 p_2}{\partial r_2^2} = 16(A+B)(\nu_{1,2} + \nu_{1,1} - 2)(6r_2^2 + 1) + \\ 16(2A+B+\Gamma)(\nu_{1,2} + \nu_{1,1} - 2)r_3^2 - 144(A+B)(\nu_{1,2} - \nu_{1,1})r_2 r_3 \\ \leq 8(A+B)(3(\nu_{1,2} + 7\nu_{1,1} - 8)r_2^2 + 2(\nu_{1,2} + \nu_{1,1} - 2)) + \\ 8(-(5A+7B-2\Gamma)\nu_{1,2} + (13A+11B+2\Gamma)\nu_{1,1} - 4(2A+B+\Gamma))r_3^2.$$

Instead of (27) and (28) we have

$$-(5A+7B-2\Gamma)\nu_{1,2} + (13A+11B+2\Gamma)\nu_{1,1} - 4(2A+B+\Gamma) \leq \\ -(5A+7B-2\Gamma)\nu_{0,2} + (13A+11B+2\Gamma)\nu_{0,1} - 4(2A+B+\Gamma) = \\ (\nu_{0,2} - \nu_{0,1}) \int_0^1 \frac{J_4}{\sqrt{1-t^2}\sqrt{1-f^2t^2}} dt$$

where

$$J_4 = -\frac{\nu_{0,2} - \nu_{0,1}}{\nu_{0,2} - 1}(3(4\nu_{0,2} - 5\nu_{0,1} + 1)t^2 + \nu_{0,2}(5\nu_{0,1} + 7) + \nu_{0,1}(-13\nu_{0,1} - 3) + 4).$$

The non-positivity of p_2 follows from

$$\begin{aligned} J_4|_{t=0} &= -13\nu_{0,1}^2 + 5\nu_{0,1}\nu_{0,2} - 3\nu_{0,1} + 7\nu_{0,2} + 4, \\ J_4|_{t=1} &= \frac{\nu_{0,1} - 1}{\nu_{0,2} - 1}(5\nu_{0,2}^2 - 13\nu_{0,1}\nu_{0,2} - 2\nu_{0,1} + 6\nu_{0,2} + 4), \end{aligned}$$

(1) and (22).

As to the coefficient p_3 , Equations (25) and (26) are replaced by

$$\begin{aligned} \frac{\partial^2 p_3}{\partial r_2^2} &= 16(\nu_{1,2} - 1)(A + B)(6r_2^2 + 1) + \\ &\quad 16((A + \Gamma)\nu_{1,2} + (A + B)\nu_{1,1} - (2A + B + \Gamma))r_3^2 - 48(\nu_{1,2} - \nu_{1,1})(A + B)r_2r_3 \\ &\quad \leq 8(A + B)(3(3\nu_{1,2} + \nu_{1,1} - 4)r_2^2 + 2(\nu_{1,2} - 1)) + \\ &\quad \quad 8(-(A + 3B - 2\Gamma)\nu_{1,2} + 5(A + B)\nu_{1,1} - 2(2A + B + \Gamma))r_3^2 \end{aligned}$$

and (27) and (28) by

$$\begin{aligned} -(A + 3B - 2\Gamma)\nu_{1,2} + 5(A + B)\nu_{1,1} - 2(2A + B + \Gamma) &\leq \\ -(A + 3B - 2\Gamma)\nu_{0,2} + 5(A + B)\nu_{0,1} - 2(2A + B + \Gamma) &= \\ (\nu_{0,2} - \nu_{0,1}) \int_0^1 \frac{J_5}{\sqrt{1-t^2}\sqrt{1-f^2t^2}} & \end{aligned}$$

where

$$J_5 = -\frac{\nu_{0,2} - \nu_{0,1}}{\nu_{0,2} - 1}((4\nu_{0,2} - 5\nu_{0,1} + 1)t^2 + \nu_{0,2}(\nu_{0,1} + 3) - \nu_{0,1}(5\nu_{0,1} + 1) + 2).$$

The non-positivity of p_3 follows from

$$\begin{aligned} J_5|_{t=0} &= -5\nu_{0,1}^2 + \nu_{0,1}\nu_{0,2} - \nu_{0,1} + 3\nu_{0,2} + 2, \\ J_5|_{t=1} &= \frac{\nu_{0,1} - 1}{\nu_{0,2} - 1}(\nu_{0,2}^2 - 5\nu_{0,1}\nu_{0,2} + 2\nu_{0,2} + 2). \end{aligned}$$

and (23), (24). □

The negativity of the only remaining Bernstein coefficient can be shown directly without resorting to Lemma 11:

Lemma 13. *The coefficient p_4 is negative.*

Proof. We can write

$$(31) \quad \frac{p_4}{16} = (\nu_{1,2} - 1)r_2^2((A + B)r_1^2 - (B - \Gamma)r_3^2) + (\nu_{1,1} - 1)r_3^2((A + \Gamma)r_1^2 + (B - \Gamma)r_2^2).$$

The proof is finished, if we can show that the coefficients of $(\nu_{1,2} - 1)r_2^2$ and $(\nu_{1,1} - 1)r_3^2$ in (31) are negative. For the coefficient of $(\nu_{1,2} - 1)r_2^2$ we argue as follows: $A < B < \Gamma < 0$ implies $A + B < A < B < B - \Gamma$ and $r_1^2 - r_2^2 - r_3^2 = 1$ implies $r_1^2 > r_3^2$. Thus, $(A + B)r_1^2 - (B - \Gamma)r_3^2 < 0$. The negativity of the coefficient of $(\nu_{1,2} - 1)r_3^2$ follows from $A + \Gamma < 0$ and $B - \Gamma < 0$. □

ACKNOWLEDGMENTS

The authors gratefully acknowledge support of this research by the Austrian Science Foundation FWF under grant P21032 (Uniqueness Results for Extremal Quadrics).

REFERENCES

- [1] K. M. Ball. Ellipsoids of maximal volume in convex bodies. *Geom. Dedicata*, 41(2):241–250, 1992.
- [2] J. Bastero and M. Romance. John’s decomposition of the identity in the non-convex case. *Positivity*, 6(1):1–16, 2002.
- [3] J. J. Callahan. *The Geometry of Spacetime. An Introduction to Special and General Relativity*. Springer, New York, 2000.
- [4] Ch. Davis. All convex invariant functions of Hermitian matrices. *Arch. Math.*, 8(4):276–278, 1957.
- [5] Y. Gordon, A. E. Litvak, M. Meyer, and A. Pajor. John’s decomposition of the identity in the general case and applications. *J. Differential Geom.*, 68(1):99–119, 2004.
- [6] P. M. Gruber. Results of baire category type in convexity. In J. Goodman, E. Lutwak, E. Malkewitsch, and J. Pollack, editors, *Discrete Geometry and Convexity*, pages 163–169. New York Academy of Sciences, 1985.
- [7] P. M. Gruber. Baire categories in convexity. In Peter M. Gruber and Jörg Wills, editors, *Handbook of convex geometry*, volume B, pages 1327–1346. Elsevier, 1993.
- [8] P. M. Gruber. Application of an idea of Voronoi to John type problems. *Adv. in Math.*, 218(2):309–351, 2008.
- [9] P. M. Gruber and F. E. Schuster. An arithmetic proof of John’s ellipsoid theorem. *Arch. Math.*, 85:82–88, 2005.
- [10] F. John. *Studies and essays. Courant anniversary volume*, chapter Extremum problems with inequalities as subsidiary conditions, pages 187–204. Interscience Publ. Inc., New York, 1948.
- [11] A. S. Lewis. Convex analysis on the Hermitian matrices. *SIAM J. Optim.*, 6(1):164–177, 1996.
- [12] E. Lutwak, D. Yang, and G. Zhang. L_p John ellipsoids. *Proc. London Math. Soc.*, 90:497–520, 2005.
- [13] M. Özdemir and A. A. Ergin. Rotations with unit timelike quaternions in Minkowski 3-space. *J. Geom. Phys.*, 56(2):322–336, 2006.
- [14] W. F. Reynold. Hyperbolic geometry on a hyperboloid. *Amer. Math. Monthly*, 100(5):442–455, 1993.
- [15] H.-P. Schröcker. Minimal enclosing hyperbolas of line sets. *Beitr. Algebra Geom.*, 48(2):367–381, 2007.
- [16] H.-P. Schröcker. Uniqueness results for minimal enclosing ellipsoids. *Comput. Aided Geom. Design*, 25(9):756–762, 2008.
- [17] M. J. Weber and H.-P. Schröcker. Davis’ convexity theorem and extremal ellipsoids. *Beitr. Algebra Geom.*, 51(1):263–274, 2010.
- [18] M. J. Weber and H.-P. Schröcker. Minimal area conics in the elliptic plane. Submitted for publication, 2010. URL <http://arxiv.org/abs/1008.4285>.

MATTHIAS. J. WEBER, HANS-PETER SCHRÖCKER, UNIT GEOMETRY AND CAD, UNIVERSITY INNSBRUCK, TECHNIKERSTRASSE 13, 6020 INNSBRUCK, AUSTRIA